

1 Assignment 1 - Solving Nonlinear Equations; Due to November 6, 2009

- The function $f(x) = 2x - \tan(x)$ is given. Solve $f(x) = 0$ in $[-1.4, 1.4]$ by:

1. Halving the Interval (Bisection) Method
2. The Method of False Position (regula falsi)
3. Newton's Method
4. Muller's Method
5. Fixed-point Iteration; $x = g(x)$ Method

- Hints:

```
>> fplot('2*x-tan(x)', [-1.4 1.4])
>> [X,FVAL]=fzero('2*x-tan(x)', [-1.4 -0.5])
>> [X,FVAL]=fzero('2*x-tan(x)', [-0.5 0.5])
>> [X,FVAL]=fzero('2*x-tan(x)', [0.5 1.4])
```

- Which methods? See Table 1

Table 1: You should use the following methods depending on the last digit of your Student ID.

Last Digit	Methods
0-1	1,2,3
2-3	1,2,4
4-5	1,2,5
6-7	1,3,4
8-9	1,3,5

- Write one complete program
 - An example calling this program;


```
>> myfunction(fx,-1.4,1.4,tol_x,tol_y)
```
 - Tabulate the actual error and function values as given in Table 2. (The number of iterations is not limited to or defined as 15. r corresponds to exact value of root.)
 - Plot the behaviours of the errors (use ratios) for the all three cases. Compare and discuss the rate of convergence.
 - You can make use of the available MATLAB codes presented in the Hands-On sessions. But, you should rearrange/rewrite all these code segments as one program (not as function calls).

n	Method1 ($x_n - r$)	Method2 ($x_n - r$)	Method3 ($x_n - r$)	n	Method1 $f(x_n)$	Method2 $f(x_n)$	Method3 $f(x_n)$
1							
2							
3							
4							
5							
6							
7							
8							
9							
10							
12							
13							
14							
15							

Table 2: Error Sequences