



**İzmir Kâtip Çelebi University**  
**Department of Engineering Sciences**  
**Phy102 Physics II**  
**Midterm Examination**  
**November 10, 2022 17:00 – 18:30**  
**Good Luck!**

**NAME-SURNAME:**

**SIGNATURE:**

**ID:**

**DEPARTMENT:**

**INSTRUCTOR:**


**DURATION:** 90 minutes

- ◇ Answer all the questions.
- ◇ Write the solutions explicitly and clearly.  
Use the physical terminology.
- ◇ You are allowed to use Formulae Sheet.
- ◇ Calculator is allowed.
- ◇ You are not allowed to use any other  
electronic equipment in the exam.

Question	Grade	Out of
1A		15
1B		15
2		20
3		20
4		20
5		20
<b>TOTAL</b>		110

This page is intentionally left blank. Use the space if needed.

1. A) In Figure, four particles form a square. The particles have charges  $q_1 = 100 \text{ nC}$ ,  $q_2 = -100 \text{ nC}$ ,  $q_3 = 200 \text{ nC}$ ,  $q_4 = -200 \text{ nC}$ , and distance  $a = 5.0 \text{ cm}$ .



i What are the  $x$  and  $y$  components of the net electrostatic force on particle 3?

ii If the charges were  $q_1 = q_4 = Q$  and  $q_2 = q_3 = q$ . What is  $Q/q$  if the net electrostatic force on particles 1 and 4 is zero?

$q_1 = 100 \times 10^{-9} \text{ C}$   
 $q_2 = -q_1$   
 $q_3 = 200 \times 10^{-9} \text{ C}$   
 $q_4 = -q_3$   
 $a = 5 \times 10^{-2} \text{ m}$

i)  $F_{3, \text{net}, x}$  &  $F_{3, \text{net}, y}$ ?  $\vec{F}_{3, \text{net}} = \sum_{i=1}^3 \vec{F}_{2i} = \vec{F}_{31} + \vec{F}_{32} + \vec{F}_{34}$  (2)

$\vec{F}_{3, \text{net}, x} = |\vec{F}_{34}| + |\vec{F}_{32}| \cos 45^\circ$  (1)  
 $\vec{F}_{3, \text{net}, y} = |\vec{F}_{32}| \sin 45^\circ - |\vec{F}_{31}|$  (1)

$F_{3, \text{net}, x} = \frac{k|q_3||q_4|}{a^2} + \frac{k|q_3||q_2|}{(a\sqrt{2})^2} \frac{\sqrt{2}}{2} = \frac{k|q_3|}{a^2} \left( |q_4| + \frac{|q_2|\sqrt{2}}{2} \right) = \frac{8.99 \times 10^9 \text{ Nm}^2/\text{C}^2 (200 \times 10^{-9} \text{ C})}{(5 \times 10^{-2} \text{ m})^2} \left( |100 \times 10^{-9} \text{ C}| + \frac{|100 \times 10^{-9} \text{ C}| \sqrt{2}}{2} \right)$   
 $= 0.169 \text{ N}$  (1) (1)

$F_{3, \text{net}, y} = \frac{k|q_3|}{a^2} \left( \frac{|q_2|\sqrt{2}}{2} - |q_1| \right) = \frac{8.99 \times 10^9 \text{ Nm}^2/\text{C}^2 (200 \times 10^{-9} \text{ C})}{(5 \times 10^{-2} \text{ m})^2} \left( \frac{100 \times 10^{-9} \text{ C} \sqrt{2}}{2} - 100 \times 10^{-9} \text{ C} \right)$   
 $= -0.046 \text{ N}$  (1) (1)

ii)  $q_1 = q_4 = Q$   
 $q_2 = q_3 = q$   
 $Q/q = ?$

$|\vec{F}_{\text{net}}| = 0 \rightarrow F_{\text{net}, x} = 0$  &  $F_{\text{net}, y} = 0$  (1)

$|\vec{F}_{\text{net}}| = 0 \rightarrow (|\vec{F}_{14}| \cos 45^\circ + |\vec{F}_{12}|)(-\hat{i}) \rightarrow (|\vec{F}_{13}| + |\vec{F}_{14}| \sin 45^\circ)(\hat{j})$

$0 = \frac{k|q_1|}{a^2} \left( \frac{|q_4|\sqrt{2}}{2} + |q_2| \right) = \frac{kQ}{a^2} \left( \frac{Q\sqrt{2}}{2} + q \right)$  (1)

$\Rightarrow \frac{Q}{q} = -\frac{4}{\sqrt{2}} = -2\sqrt{2} = -2.83$  (1)

- B) The density of conduction electrons in aluminum is  $2.1 \times 10^{29} \text{ m}^{-3}$ . What is the drift velocity in an aluminum conductor that has a  $2.0 \mu\text{m}$  by  $3.0 \mu\text{m}$  rectangular cross section and when a  $32.0 \text{ mA}$  current flows through the conductor?

$$\begin{aligned}
 n &= 2.1 \times 10^{29} \text{ m}^{-3} \\
 i &= 32 \times 10^{-3} \text{ A} \\
 A &= (2 \times 10^{-6} \text{ m})(3 \times 10^{-6} \text{ m}) \\
 v_d &=?
 \end{aligned}$$

$$\vec{J} = ne\vec{v}_d$$

$$\begin{aligned}
 \textcircled{3} J &= nev_d \\
 \textcircled{3} J &= \frac{i}{A}
 \end{aligned}
 \left. \vphantom{\begin{aligned} \textcircled{3} J &= nev_d \\ \textcircled{3} J &= \frac{i}{A} \end{aligned}} \right\} \frac{i}{A} = nev_d$$

$$\Rightarrow v_d = \frac{i}{Ane} = \frac{32 \times 10^{-3} \text{ A}}{(2 \times 10^{-6} \text{ m})(3 \times 10^{-6} \text{ m})(2.1 \times 10^{29} \text{ m}^{-3})(1.602 \times 10^{-19} \text{ C})}$$

$$= \boxed{0.016 \text{ m/s}}$$

$$\begin{aligned}
 \frac{\text{A}}{\text{m}^2 \text{ m}^{-3} \text{ C}} &\sim \frac{\text{C/s}}{\text{m}^2 \text{ m}^{-3} \text{ C}} \sim \text{m/s} \\
 &\text{unit check}
 \end{aligned}$$

2. At some instant the velocity components of an electron moving between two charged parallel plates are  $v_x = 3 \times 10^5 \text{ m/s}$  and  $v_y = 5.0 \times 10^3 \text{ m/s}$ . Suppose the electric field between the plates is given by  $\vec{E} = (180 \text{ N/C})\hat{j}$ . In unit-vector notation, what are

- the electron's acceleration in that field
- the electron's velocity when its x coordinate has changed by 2.4 cm?


$\vec{e}$ : electron  
 $v_x = 3 \times 10^5 \text{ m/s} = v_{0x}$   
 $v_y = 5 \times 10^3 \text{ m/s} = v_{0y}$   
 $\vec{E} = 180 \text{ N/C} \hat{j}$

i)  $a = ?$   
 $\vec{F}_E = q\vec{E} = (1.6 \times 10^{-19} \text{ C})(180 \text{ N/C})(-\hat{j})$   
 $= 288 \times 10^{-19} \text{ N}(-\hat{j})$   
 $\Rightarrow m_e \vec{a} = \vec{F}_E \Rightarrow \vec{a} = \frac{288 \times 10^{-19} \text{ N}(-\hat{j})}{9.109 \times 10^{-31} \text{ kg}} = \boxed{3.16 \times 10^{13} \text{ m/s}^2(-\hat{j})}$

ii)  $\Delta x = x - x_0 = 2.4 \times 10^{-2} \text{ m}$  & no force acting on x direction  
 $\Rightarrow v_x = v_{0x}$  &  $v_y = v_{0y} + at$   
 $\Rightarrow \Delta t = \frac{2.4 \times 10^{-2} \text{ m}}{3 \times 10^5 \text{ m/s}} = 8 \times 10^{-8} \text{ s}$   
 $\Rightarrow v_y = 5 \times 10^3 \text{ m/s} - (3.16 \times 10^{13} \text{ m/s}^2)(8 \times 10^{-8} \text{ s}) = \boxed{-2.52 \times 10^6 \text{ m/s}}$

$\Rightarrow \vec{v} = 3 \times 10^5 \text{ m/s} \hat{i} + 2.52 \times 10^6 \text{ m/s}(-\hat{j})$

3. A small, nonconducting ball of mass  $m = 2 \times 10^{-6} \text{ kg}$  and charge  $q = 4.0 \times 10^{-8} \text{ C}$  (distributed uniformly through its volume) hangs from an insulating thread that makes an angle  $\theta = 60^\circ$  with a vertical, uniformly charged **nonconducting sheet** (shown in cross section).



Considering the gravitational force on the ball and assuming the sheet extends far vertically and into and out of the page, **calculate the surface charge density  $\sigma$**  of the sheet. (Hint: The ball is in equilibrium (stationary).)

$m = 2 \times 10^{-6} \text{ kg}$  (non-conducting)  
 $q = 4 \times 10^{-8} \text{ C}$  uniform distribution  
 non-conducting sheet,  $\sigma = ?$  (if hanged)  
 $E = \frac{\sigma}{2\epsilon_0}$


$T \cos 60^\circ - mg = ma_y = 0$   
 $qE - T \sin 60^\circ = ma_x = 0$   
 $mg = F_g$  hangs  $\rightarrow$  stationary

now, eliminate  $T$

$\rightarrow qE - \left(\frac{mg}{\cos 60^\circ}\right) \sin 60^\circ = 0 \rightarrow qE = mg \tan 60^\circ \rightarrow q \frac{\sigma}{2\epsilon_0} = mg \tan 60^\circ \rightarrow \sigma = \frac{2mg \epsilon_0 \tan 60^\circ}{q}$

$\rightarrow \sigma = \frac{2mg \epsilon_0 \tan 60^\circ}{q} = \frac{2(2 \times 10^{-6} \text{ kg})(9.8 \text{ m/s}^2)(8.85 \times 10^{-12} \text{ C}^2/\text{Nm}^2) \tan 60^\circ}{(4 \times 10^{-8} \text{ C})} = 15 \times 10^9 \text{ C/m}^2$

4. Two very thin non-conducting rods are placed together as shown. Both rods have lengths of  $L$  and they carry uniform charges of  $+q$  and  $-q$  over their lengths. Find the potential at point  $P$  at a distance  $a$  and  $d$  from the positively and negatively charged rods as shown. Don't perform integration.




$V_{1 \text{ at } P} = \int dV$   
 $= \int \frac{1}{4\pi\epsilon_0} \frac{dq_1}{r_1}$   
 $dq_1 = \lambda dx$  (3)  
 $r_1 = L + a - x$

$V_{2 \text{ at } P} = \int dV$   
 $= \int \frac{1}{4\pi\epsilon_0} \frac{dq_2}{r_2}$   
 $dq_2 = -\lambda dx$  (3)  
 $r_2 = \sqrt{x^2 + d^2}$

$V = \frac{1}{4\pi\epsilon_0} \frac{q}{r}$   
 $dV = \frac{1}{4\pi\epsilon_0} \frac{dq}{r}$  (2)  
 $dq = ?$   
 $dq = \lambda dx$  (2)

$V_{\text{tot}} = V_{1 \text{ at } P} + V_{2 \text{ at } P} = \frac{\lambda}{4\pi\epsilon_0} \left( \int_0^L \frac{dx}{L+a-x} - \int_0^L \frac{dx}{\sqrt{x^2 + d^2}} \right)$  (3)

5. The parallel plate capacitors in the given circuit have the same plate area  $A$  and plate separation  $d$ . The capacitance of the air-filled capacitor is  $C_1 = 6.0 \mu F$ . A dielectric slab of dielectric constant  $\kappa = 2.0$  is placed between the plates of the second capacitor as shown. The voltage across the combination of capacitors is  $\Delta V = 20 V$  and the capacitors are fully charged.



- Find the equivalent capacitance of the combination of capacitors.
- Calculate the energy stored in each capacitor.
- Calculate the electric field in the second capacitor if the area of the capacitor is  $100 \text{ cm}^2$ .

i) Capacitors  $C_1$  &  $C_2$  are in series.  $C = \kappa \frac{\epsilon_0 A}{d}$  (1)

$C_1$ : air filled  $\rightarrow \kappa = 1 \Rightarrow C_1 = \frac{\epsilon_0 A}{d} = 6.0 \mu F$  (1)


$C_2$ : dielectric slab  $\rightarrow \kappa = 2 \Rightarrow C_2 = \kappa C_1 = 12.0 \mu F$  (1)

$\Rightarrow \frac{1}{C_{\text{equiv}}} = \frac{1}{C_1} + \frac{1}{C_2} \Rightarrow \frac{1}{C_{\text{equiv}}} = \frac{1}{6 \mu F} + \frac{1}{12 \mu F} = \frac{18}{72 \mu F} \Rightarrow C_{\text{equiv}} = 4 \mu F$  (1) (1)

ii) In series  $\rightarrow$  same charge on both capacitors (1)

(1)  $q = q_1 = q_2$  &  $\Delta V = 20 V$   $\left\{ \begin{array}{l} C = \frac{q}{\Delta V} \rightarrow q = C \Delta V = (4 \mu F)(20 V) = 80 \mu C \end{array} \right.$  (1)

$U = \frac{q^2}{2C} \left\{ \begin{array}{l} U_1 = \frac{(80 \mu C)^2}{2(6.0 \mu F)} = 533.3 \mu J \\ U_2 = \frac{(80 \mu C)^2}{2(12 \mu F)} = 266.6 \mu J \end{array} \right.$  (1) (1)

iii)  (1)

$A = 100 \text{ cm}^2 = 10^{-2} \text{ m}^2$  (1)

$\epsilon_0 \kappa E \cdot d \vec{A} = q$  (1)

$\Rightarrow E = \frac{q}{\epsilon_0 \kappa A} = \frac{80 \times 10^{-6} C}{(8.85 \times 10^{-12} \frac{F}{m})(2.0)(10^{-2} \text{ m}^2)} = 4.52 \times 10^3 \text{ V/m}$  (1) (1)





**İzmir Kâtip Çelebi University**  
**Department of Engineering Sciences**  
**Phy102 Physics II**  
**Midterm Examination**  
**November 11, 2021 17:00 – 18:30**  
**Good Luck!**

**NAME-SURNAME:**

**SIGNATURE:**

**ID:**

**DEPARTMENT:**

**INSTRUCTOR:**


**DURATION:** 90 minutes

- ◇ Answer all the questions.
- ◇ Write the solutions explicitly and clearly.  
Use the physical terminology.
- ◇ You are allowed to use Formulae Sheet.
- ◇ Calculator is allowed.
- ◇ You are not allowed to use any other electronic equipment in the exam.

Question	Grade	Out of
1A		15
1B		15
2		20
3		20
4		20
5		20
<b>TOTAL</b>		110

This page is intentionally left blank. Use the space if needed.

1. A) In figure, all four particles are fixed in the  $xy$ -plane, and  $q_1 = -3.20 \times 10^{-19} \text{ C}$ ,  $q_2 = +3.20 \times 10^{-19} \text{ C}$ ,  $q_3 = +6.40 \times 10^{-19} \text{ C}$ ,  $q_4 = +3.20 \times 10^{-19} \text{ C}$ ,  $\theta_1 = 35.0^\circ$ ,  $d_1 = 3.00 \text{ cm}$  and  $d_2 = d_3 = 2.00 \text{ cm}$ .



What are the magnitude and direction of the net electrostatic force on particle 4 due to the other three particles?

Target particle: 4

$q_1 = -3.2 \times 10^{-19} \text{ C}$   
 $q_2 = +3.2 \times 10^{-19} \text{ C}$   
 $q_3 = +6.4 \times 10^{-19} \text{ C}$   
 $q_4 = +3.2 \times 10^{-19} \text{ C}$   
 $\theta_1 = 35^\circ$   
 $d_1 = 3 \times 10^{-2} \text{ m}$   
 $d_2 = d_3 = 2 \times 10^{-2} \text{ m}$

$\vec{F}_{4net} = \sum_{i=1}^3 \vec{F}_{4i}$   
 $= \vec{F}_{41} + \vec{F}_{42} + \vec{F}_{43}$

$F_{4net,x} = F_{43,x} + F_{41,x}$   
 $F_{4net,y} = F_{42,y} + F_{41,y}$

$\Rightarrow F_{4net,x} = -F_{43} - F_{41} \cos 35^\circ$   
 $= -k \frac{q_4 |q_3|}{d_3^2} - k \frac{q_4 |q_1|}{d_1^2} \cos 35^\circ$   
 $= -(8.99 \times 10^9 \text{ Nm}^2/\text{C}^2) (3.2 \times 10^{-19} \text{ C})^2 \left( \frac{2}{(2 \times 10^{-2} \text{ m})^2} + \frac{\cos 35^\circ}{(3 \times 10^{-2} \text{ m})^2} \right) = -5.44 \times 10^{-24} \text{ N}$

$F_{4net,y} = -F_{42} - F_{41} \sin 35^\circ$   
 $= -k \frac{q_4 |q_2|}{d_2^2} - k \frac{q_4 |q_1|}{d_1^2} \sin 35^\circ$   
 $= -(8.99 \times 10^9 \text{ Nm}^2/\text{C}^2) (3.2 \times 10^{-19} \text{ C})^2 \left( \frac{1}{(2 \times 10^{-2} \text{ m})^2} + \frac{\sin 35^\circ}{(3 \times 10^{-2} \text{ m})^2} \right) = -2.89 \times 10^{-24} \text{ N}$

$F_{4net} = \sqrt{F_{4net,x}^2 + F_{4net,y}^2} = \sqrt{(-5.44 \times 10^{-24} \text{ N})^2 + (-2.89 \times 10^{-24} \text{ N})^2} = 6.16 \times 10^{-24} \text{ N}$

$\tan \theta = \frac{F_{4net,y}}{F_{4net,x}} = \frac{-2.89 \times 10^{-24}}{-5.44 \times 10^{-24}} \Rightarrow \theta = \tan^{-1} \frac{-2.89}{-5.44} = 27.98^\circ \approx 28^\circ$

III. quadrant  $\theta = 203^\circ$   
 magnitude  $\theta = 28^\circ$   
 angle

- B) The density of conduction electrons in aluminum is  $2.1 \times 10^{29} \text{ m}^{-3}$ . What is the drift velocity in an aluminum conductor that has a  $2.0 \mu\text{m}$  by  $3.0 \mu\text{m}$  rectangular cross section and when a  $32.0 \text{ mA}$  current flows through the conductor?

$$\begin{aligned}
 n &= 2.1 \times 10^{29} \text{ m}^{-3} \\
 i &= 32 \times 10^{-3} \text{ A} \\
 A &= (2 \times 10^{-6} \text{ m})(3 \times 10^{-6} \text{ m}) \\
 v_d &=?
 \end{aligned}$$

$$\begin{aligned}
 \vec{J} &= ne\vec{v}_d \\
 \textcircled{3} J &= nev_d \\
 \textcircled{3} J &= \frac{i}{A}
 \end{aligned}
 \quad \left. \vphantom{\begin{aligned} \vec{J} &= nev_d \\ \textcircled{3} J &= nev_d \\ \textcircled{3} J &= \frac{i}{A} \end{aligned}} \right\} \frac{i}{A} = nev_d$$


$$\Rightarrow v_d = \frac{i}{Ane} = \frac{32 \times 10^{-3} \text{ A}}{(2 \times 10^{-6} \text{ m})(3 \times 10^{-6} \text{ m})(2.1 \times 10^{29} \text{ m}^{-3})(1.602 \times 10^{-19} \text{ C})}$$

$$= \boxed{0.016 \text{ m/s}}$$

$$\begin{aligned}
 \frac{\text{A}}{\text{m}^2 \text{ m}^{-3} \text{ C}} &\sim \frac{\text{C/s}}{\text{m}^2 \text{ m}^{-3} \text{ C}} \sim \text{m/s} \\
 &\text{unit check}
 \end{aligned}$$

2. Semicircular wire shown in figure below has a non-uniform charge distribution  $\lambda(\theta) = \lambda_0 \cos\theta$ .

Find the electric field at point P in unit vector notation and in terms of total charge Q.  
 (Hint:  $\int \cos^2 ax dx = x/2 + \sin 2ax/4a$ )



$$dE = \frac{k dq}{R^2} = \frac{k \lambda R d\theta}{R^2} = \frac{k \lambda_0 \cos\theta}{R} d\theta$$

$$x\text{-components are cancelling due to symmetry}$$

$$dE_y = |dE| \cos\theta = \frac{k \lambda_0 \cos^2\theta}{R} d\theta$$

$$E_y = E = \frac{k \lambda_0}{R} \int_{-\pi/2}^{\pi/2} \cos^2\theta d\theta = \frac{k \lambda_0}{R} \left[ \frac{\theta}{2} + \frac{\sin 2\theta}{4} \right]_{-\pi/2}^{\pi/2}$$


$$= \frac{k \lambda_0 \pi}{2R} \Rightarrow \vec{E} = \frac{k \lambda_0 \pi}{2R} \hat{j}$$

in terms of Q

$$Q = \int \lambda ds = \int_{-\pi/2}^{\pi/2} \lambda_0 \cos\theta R d\theta = \lambda_0 R \sin\theta \Big|_{-\pi/2}^{\pi/2} = 2\lambda_0 R$$

$$\Rightarrow \vec{E} = \frac{1}{4\pi\epsilon_0} \frac{Q}{2R} \frac{\pi}{2R} \hat{j} = \frac{Q}{16\pi\epsilon_0 R^2} \hat{j}$$

3. Figure shows a spherical shell with uniform volume charge density  $\rho = (1.56 \times 10^{-9} \text{ C/m}^3)$ , inner radius  $a = 10 \text{ cm}$ , and outer radius  $b = 2.00a$ .




What is the magnitude of the electric field at radial distances

i  $r = 1.5a$

ii  $r = 3.00b$

Hints: Use Gauss' Law. Volume of the spherical shell:  $\frac{4}{3}\pi(b^3 - a^3)$ .



$\rho = 1.56 \times 10^{-9} \text{ C/m}^3$   
 $a = 10 \times 10^{-2} \text{ m}$   
 $b = 2a$   
 $r_i = 1.5a$   
 $r_u = 3b$

$V_{ss} = \frac{4}{3}\pi(b^3 - a^3)$

$\oint \vec{E} \cdot d\vec{A} = \frac{q_{enc}}{\epsilon_0}$  ,  $\rho = \frac{q}{V} = \frac{q_{enc}}{V_{enc}}$  ,  $\vec{E} \parallel \vec{A}$   
 i)  $\oint \vec{E} \cdot d\vec{A} = \frac{q_{enc}}{\epsilon_0} \Rightarrow E 4\pi r_i^2 = \frac{q_{enc}}{\epsilon_0}$  }  $q_{enc} = ?$   
 $G S_i \Rightarrow q_{enc} = \rho V_{enc} = \rho \frac{4}{3}\pi(r_i^3 - a^3)$   
 $\Rightarrow E 4\pi r_i^2 = \rho \frac{4}{3}\pi(r_i^3 - a^3)$  }  $r_i = 1.5a$   
 $\Rightarrow E = \frac{\rho \frac{4}{3}\pi}{4\pi\epsilon_0} \frac{(1.5a)^3 - a^3}{1.5a^2} = \frac{\rho a}{3\epsilon_0} \left(\frac{2.375}{2.25}\right)$   
 $E(r=1.5a) = \frac{(1.56 \times 10^{-9} \text{ C/m}^3)(10 \times 10^{-2} \text{ m})}{3 \times 8.85 \times 10^{-12} \text{ C}^2/(\text{Nm}^2)} \left(\frac{2.375}{2.25}\right) = \boxed{6.20 \text{ N/C}}$   
 ii)  $E 4\pi r_u^2 = \frac{q_{enc}}{\epsilon_0} \Rightarrow q_{enc} = \rho \frac{4}{3}\pi(b^3 - a^3)$   
 $G S_u \Rightarrow E 4\pi r_u^2 = \rho \frac{4}{3}\pi(b^3 - a^3)$  }  $r_u = 3b$   
 $\Rightarrow E = \frac{\rho \frac{4}{3}\pi}{4\pi\epsilon_0} \frac{b^3 - a^3}{(3b)^2} = \frac{\rho 7a}{108\epsilon_0}$   
 $E(r=3b) = \frac{(1.56 \times 10^{-9} \text{ C/m}^3) 7(10 \times 10^{-2} \text{ m})}{108 \times 8.85 \times 10^{-12} \text{ C}^2/(\text{Nm}^2)} = \boxed{1.14 \text{ N/C}}$

4. The electric potential at points in an xy plane is given by  $V = 4x^2 - 2y^3$ .  
**In unit vector notations**, what is the electric field at point (1m, 2m)?

$$V(x,y) = 4x^2 - 2y^3 \quad \& \quad E_s = -\frac{\partial V}{\partial s}$$

$$\vec{E} = -\frac{\partial V}{\partial x} \hat{i} - \frac{\partial V}{\partial y} \hat{j} = -8x \hat{i} + 6y^2 \hat{j}$$

$$\vec{E}(x=1\text{m}, y=2\text{m}) = \boxed{-8 \hat{i} + 24 \hat{j}}$$

5. In figure below, the parallel plate capacitor of plate area  $2 \times 10^{-2} \text{ m}^2$  is filled with two dielectric slabs, each with thickness  $2.00 \text{ mm}$ . One slab has dielectric constant 3.00, and the other, 4.00. **How much charge** does the  $7.00 \text{ V}$  battery store on the capacitor?



$A = 2 \times 10^{-2} \text{ m}^2$   
 $d = 2 \times 10^{-3} \text{ m}$   
 $K_1 = 3 \text{ \& } K_2 = 4$   
 $V = 7 \text{ V}$   
 $q = ?$

in series connection

$C_1 = K_1 \epsilon_0 \frac{A}{d} = 3 \epsilon_0 \frac{A}{d}$   
 $C_2 = K_2 \epsilon_0 \frac{A}{d} = 4 \epsilon_0 \frac{A}{d}$

$C_{\text{equiv}} = \frac{C_1 C_2}{C_1 + C_2} = \frac{12}{7} \epsilon_0 \frac{A}{d} = \frac{12}{7} (8.85 \times 10^{-12} \text{ C}^2 / (\text{Nm}^2)) \frac{2 \times 10^{-2} \text{ m}^2}{2 \times 10^{-3} \text{ m}}$   
 $= 1.52 \times 10^{-10} \text{ F}$

$C_{\text{equiv}} = \frac{Q}{V} \Rightarrow q = C_{\text{equiv}} V = (1.52 \times 10^{-10} \text{ F}) 7 \text{ V} = 1.06 \times 10^{-9} \text{ C}$

$\frac{\text{C}^2 \text{ m}^2}{\text{Nm}^2 \text{ m}} \sim \text{F} \sim \frac{\text{C}}{\text{V}} = \frac{\text{C}}{\text{J/C}} = \frac{\text{C}^2}{\text{J}}$   
 mit check





**İzmir Kâtip Çelebi University**  
**Department of Engineering Sciences**  
**Phy102 Physics II**  
**Midterm Examination**  
**November 03, 2019 15:30 – 17:30**  
**Good Luck!**

**NAME-SURNAME:**

**SIGNATURE:**

**ID:**

**DEPARTMENT:**

**INSTRUCTOR:**

**DURATION:** 120 minutes

- ◇ Answer all the questions.
- ◇ Write the solutions explicitly and clearly.  
Use the physical terminology.
- ◇ You are allowed to use Formulae Sheet.
- ◇ Calculator is allowed.
- ◇ You are not allowed to use any other electronic equipment in the exam.

Question	Grade	Out of
1A		15
1B		15
2		20
3		20
4		20
5		20
<b>TOTAL</b>		110

This page is intentionally left blank. Use the space if needed.

1. A) A point charge  $q_1 = 8 \text{ nC}$  is at the origin and a second point charge  $q_2 = 12 \text{ nC}$  is on the x-axis at  $x=4 \text{ m}$ . Find the net electric force they exert on  $q_3 = -5 \text{ nC}$  located on the y-axis at  $y=3.0 \text{ m}$  in vector notation, magnitude and angle.

$q_3 = -5 \text{ nC}$   
 $q_1 = 8 \text{ nC}$   
 $q_2 = 12 \text{ nC}$

$\vec{F}_{3, \text{net}} = \vec{F}_{31} + \vec{F}_{32}$

$|\vec{F}_{31}| = k \frac{|q_3| |q_1|}{r_{31}^2} = \left(9 \times 10^9 \frac{\text{Nm}^2}{\text{C}^2}\right) \frac{5 \times 10^{-9} \text{C} |8 \times 10^{-9} \text{C}|}{(3 \text{ m})^2}$   
 $= 4 \times 10^{-8} \text{ N} \rightarrow \vec{F}_{31} = 4 \times 10^{-8} \text{ N} (\hat{j})$

$|\vec{F}_{32}| = k \frac{|q_3| |q_2|}{r_{32}^2} = \left(9 \times 10^9 \frac{\text{Nm}^2}{\text{C}^2}\right) \frac{5 \times 10^{-9} \text{C} |12 \times 10^{-9} \text{C}|}{(4 \text{ m})^2}$   
 $= 2.16 \times 10^{-8} \text{ N} \rightarrow \vec{F}_{32} = ?$

$\cos \theta = \frac{4}{5} = 0.8$   
 $\sin \theta = \frac{3}{5} = 0.6$


$F_{32, x} = |\vec{F}_{32}| \cos \theta = 2.16 \times 10^{-8} \text{ N} \times 0.8 = 1.73 \times 10^{-8} \text{ N}$   
 $F_{32, y} = |\vec{F}_{32}| \sin \theta = 2.16 \times 10^{-8} \text{ N} \times 0.6 = 1.3 \times 10^{-8} \text{ N}$

$\Rightarrow \vec{F}_{3, \text{net}} = (4 \times 10^{-8} \hat{j}) + (1.73 \times 10^{-8} \hat{i} + 1.3 \times 10^{-8} \hat{j}) \text{ N} = 1.73 \times 10^{-8} \hat{i} - 5.3 \times 10^{-8} \hat{j}$

$|\vec{F}_{3, \text{net}}| = \sqrt{(1.73 \times 10^{-8} \text{ N})^2 + (-5.3 \times 10^{-8} \text{ N})^2} = 5.6 \times 10^{-8} \text{ N}$   
 $\theta = \tan^{-1} \frac{-5.3 \text{ N}}{1.73} = -72^\circ$

- B) Semicircular wire shown in figure below has a non-uniform charge distribution  $\lambda(\theta) = \lambda_0 \cos\theta$ .

Find the electric field at point P in unit vector notation and in terms of total charge Q.  
 (Hint:  $\int \cos^2 ax dx = x/2 + \sin 2ax/4a$ )



$$dE = \frac{k dq}{R^2} = \frac{k \lambda R d\theta}{R^2} = \frac{k \lambda_0 \cos\theta}{R} d\theta$$

$$x\text{-components are cancelling due to symmetry}$$

$$dE_y = dE \cos\theta = \frac{k \lambda_0 \cos^2\theta}{R} d\theta$$

$$E_y = E = \frac{k \lambda_0}{R} \int_{-\pi/2}^{\pi/2} \cos^2\theta d\theta = \frac{k \lambda_0}{R} \left[ \frac{\theta}{2} + \frac{\sin 2\theta}{4} \right]_{-\pi/2}^{\pi/2}$$

$$= \frac{k \lambda_0 \pi}{2R} \Rightarrow \vec{E} = \frac{k \lambda_0 \pi}{2R} \hat{j}$$

in terms of Q


$$Q = \int \lambda ds = \int_{-\pi/2}^{\pi/2} \lambda_0 \cos\theta R d\theta = \lambda_0 R \sin\theta \Big|_{-\pi/2}^{\pi/2} = 2\lambda_0 R$$

$$\Rightarrow \vec{E} = \frac{1}{4\pi\epsilon_0} \frac{Q}{2R} \frac{\pi}{2R} \hat{j} = \frac{Q}{16\pi\epsilon_0 R^2} \hat{j}$$

2. A proton moves at  $4.5 \times 10^5 \text{ m/s}$  in the horizontal direction. It enters a uniform vertical electric field with a magnitude of  $9.6 \times 10^3 \text{ N/C}$ .

Ignoring any gravitational effects, find

- the time required for the proton to travel 5 cm horizontally,
- the vertical displacement during that time,
- the horizontal and vertical components of the velocity after the proton has traveled 5 cm horizontally.



$v = 4.5 \times 10^5 \text{ m/s}$  &  $E = 9.6 \times 10^3 \text{ N/C}$   
 (uniform)  $\rightarrow$   
 Constant  $E \rightarrow$  constant acceleration  $\leftarrow$  force  
 $v = v_{0x}$  &  $v_{0y} = 0$   
 $a = a_y$  &  $a_x = 0$

$qE = ma$


i)  $v_x = v_{0x} = v = \frac{\Delta x}{\Delta t} \rightarrow \Delta t = \frac{5 \times 10^{-2} \text{ m}}{4.5 \times 10^5 \text{ m/s}} = 1.11 \times 10^{-7} \text{ s} = \underline{\underline{111 \text{ ns}}}$

ii)  $a_y m_p = q_p E \rightarrow a_y = \frac{q_p E}{m_p} = \frac{(1.6 \times 10^{-19} \text{ C})(9.6 \times 10^3 \text{ N/C})}{(1.67 \times 10^{-27} \text{ kg})} = 9.21 \times 10^{10} \text{ m/s}^2$

$y = y_0 + v_{0y} t + \frac{1}{2} a_y t^2 \rightarrow y - y_0 = h = \frac{1}{2} a_y t^2 = \frac{1}{2} (9.21 \times 10^{10} \text{ m/s}^2) (1.11 \times 10^{-7} \text{ s})^2$   
 $= 5.68 \times 10^{-3} \text{ m} = \underline{\underline{5.68 \text{ mm}}}$

iii)  $v_x = v_{0x} = 4.5 \times 10^5 \text{ m/s}$   
 $v_y = v_{0y} + a_y t = a_y t = (9.21 \times 10^{10} \text{ m/s}^2) (1.11 \times 10^{-7} \text{ s}) = \underline{\underline{1.02 \times 10^5 \text{ m/s}}}$

3. Figure below shows a section of a conducting rod of radius  $R_1 = 1.30 \text{ mm}$  and length  $L = 11.00 \text{ m}$  inside a thin-walled coaxial conducting cylindrical shell of radius  $R_2 = 10.0R_1$  and the (same) length  $L$ . The net charge on the rod is  $Q_1 = +3.40 \times 10^{-12} \text{ C}$ ; that on the shell is  $Q_2 = -2.00Q_1$



- What are the magnitude  $E$  and direction (radially inward or outward) of the electric field at radial distance  $r = 2.00R_2$ ?
- What are  $E$  and the direction at  $r = 5.00R_1$ ?
- What is the charge on the interior and exterior surface of the shell?

$R_1 = 1.30 \times 10^{-3} \text{ m}$   
 $R_2 = 10.0 R_1 = 1.30 \times 10^{-2} \text{ m}$   
 $L = 11.00 \text{ m}$

$Q_1 = +3.40 \times 10^{-12} \text{ C}$  (on rod)  
 $Q_2 = -2 Q_1 = -6.80 \times 10^{-12} \text{ C}$  (on shell)


$\oint \vec{E} \cdot d\vec{A} = \frac{q_{\text{enc}}}{\epsilon_0}$  (cylindrical Gaussian surface)

i) GS1:  $r = 2R_2 \rightarrow E 2\pi r L = \frac{Q_1 + Q_2}{\epsilon_0} \Rightarrow E = \frac{3.4 \times 10^{-12} - 6.8 \times 10^{-12}}{2\pi \times 1.30 \times 10^{-2} \text{ m} \times 11 \text{ m} \times \epsilon_0}$   
 $\vec{E} \rightarrow \hat{r}_1 \rightarrow E = -0.214 \text{ N/C} \rightarrow |\vec{E}| = 0.214 \text{ N/C}$  & inward

ii) GS2:  $r = 5R_1 \rightarrow E 2\pi r L = \frac{Q_1}{\epsilon_0} \rightarrow E = \frac{3.4 \times 10^{-12}}{2\pi \times 1.30 \times 10^{-3} \text{ m} \times 11 \text{ m} \times \epsilon_0} = 8.55 \times 10^{-12} \text{ C}^2/\text{Nm}^2$   
 $\vec{E} \rightarrow \hat{r}_2 \rightarrow E = 0.855 \text{ N/C} \rightarrow |\vec{E}| = 0.855 \text{ N/C}$  & outward

iii)  $Q_1$  (rod outer)  $-Q_1$  (shell inner)  $Q_2 - (-Q_1) \rightarrow 3.4 \times 10^{-12} - 3.4 \times 10^{-12} - 6.8 \times 10^{-12} - (-3.4 \times 10^{-12})$   
 sum up to  $-6.8 \times 10^{-12} \text{ C}$

4. An electric dipole of two opposite charges of magnitude  $q = 1.50 \mu\text{C}$ , separated by a distance  $d = 1.20 \text{ cm}$  is placed near an infinitely large plane of charge of uniform charge density  $\sigma = 1.77 \mu\text{C}/\text{m}^2$ . The axis of the electric dipole makes an angle of  $\varphi = 37^\circ$  with the plane, as shown in the figure.



- i Find the magnitude of the electric field due to the plane. Show its direction on the figure.
- ii Calculate the magnitude of the electric dipole moment. Show its direction on the figure.
- iii Calculate the magnitude of the torque acting on the electric dipole. Show its direction on the figure.
- iv How much work must be done by an external agent to turn the electric dipole by  $90^\circ$  in clockwise direction?

$\vec{E}$   
 $\vec{E}$   
 $\vec{E}$   
 $\vec{E}$   
 $\vec{E}$   
 $\vec{E}$   
 $\vec{E}$   
 $\vec{E}$

i)  $\sigma = 1.77 \times 10^{-6} \text{ C/m}^2$  & non-conducting plane (uniform charge density)

$$\Rightarrow |\vec{E}| = \frac{\sigma}{2\epsilon_0} = \frac{1.77 \times 10^{-6} \text{ C/m}^2}{2 \times 8.85 \times 10^{-12} \text{ C}^2/\text{Nm}^2} = 10^5 \text{ N/C}$$

magnitude

ii)  $p = qd$   $\left\{ \begin{array}{l} \ominus \text{---} d \text{---} \oplus \\ \vec{p} \end{array} \right.$   $p = 1.50 \times 10^{-6} \text{ C} \times 1.20 \times 10^{-2} \text{ m} = 1.8 \times 10^{-8} \text{ Cm}$

iii)  $\vec{\tau} = \vec{p} \times \vec{E} \Rightarrow |\vec{\tau}| = |\vec{p}| |\vec{E}| \sin 53^\circ = (1.8 \times 10^{-8} \text{ Cm})(10^5 \text{ N/C}) \sin 53^\circ$

$$= 1.44 \times 10^{-3} \text{ Nm} \quad \otimes \text{ into the page}$$

iv) rotation in cw,  $90^\circ$

initial,  $\vec{p}_i$   
 final,  $\vec{p}_f$

$$W_{\text{ext}} = -\Delta U = -U_f + U_i = U_i - U_f \quad \left\{ U = \vec{p} \cdot \vec{E} \right.$$


$$= \vec{p}_i \cdot \vec{E} - \vec{p}_f \cdot \vec{E}$$

$$= |\vec{p}| |\vec{E}| \cos 53^\circ - |\vec{p}| |\vec{E}| \cos 37^\circ$$

$$= (1.8 \times 10^{-8} \text{ Cm})(10^5 \text{ N/C})(\cos 53^\circ - \cos 37^\circ)$$

$$= -3.6 \times 10^{-4} \text{ J} \quad \text{J} \equiv \text{Nm}$$

5. Four capacitors are connected as shown in Figure.



- i Find the equivalent capacitance between points a and b.
- ii Calculate the charge on each capacitor if  $\Delta V_{ab} = 15.0 \text{ V}$ .

i)  $C_{12} = \frac{C_1 C_2}{C_1 + C_2} = \frac{(15 \times 10^{-6} \text{ F})(3 \times 10^{-6} \text{ F})}{18 \times 10^{-6} \text{ F}} = 2.5 \times 10^{-6} \text{ F}$   
 $C_{123} = C_{12} + C_3 = 2.5 \times 10^{-6} \text{ F} + 6 \times 10^{-6} \text{ F} = 8.5 \times 10^{-6} \text{ F}$   
 $C_{eq} = C_{1234} = \frac{C_{123} C_4}{C_{123} + C_4} = \frac{(8.5 \times 10^{-6} \text{ F})(20 \times 10^{-6} \text{ F})}{28.5 \times 10^{-6} \text{ F}} = 5.97 \times 10^{-6} \text{ F} = 5.97 \mu\text{F}$

ii)  $C = \frac{Q}{V} \rightarrow Q_{eq} = Q_{1234} = C_{eq} V = 5.97 \times 10^{-6} \text{ F} \times 15 \text{ V} = 89.47 \mu\text{C}$   
 $\rightarrow Q_4 = Q_{123} = Q_{eq} = 89.47 \mu\text{C} \rightarrow V_4 = \frac{89.47 \mu\text{C}}{20 \mu\text{F}} = 4.47 \text{ V} \leftarrow (4)$

$10.53 \text{ V} \leftarrow (3) \rightarrow Q_3 = C_3 V_3 = 63.18 \mu\text{C} \leftarrow (3)$   
 $Q_{12} = C_{12} V = (2.5 \times 10^{-6} \text{ F}) 10.53 \text{ V} = Q_1 = Q_2$   
 $\rightarrow Q_1 = Q_2 = 2.63 \mu\text{C}$

$\Rightarrow V_1 = \frac{Q_1}{C_1} = \frac{2.63 \mu\text{C}}{15 \mu\text{F}} = 1.75 \text{ V} \leftarrow (1)$   
 $V_2 = \frac{Q_1}{C_2} = 8.78 \text{ V} \leftarrow (2)$





**İzmir Kâtip Çelebi University**  
**Department of Engineering Sciences**  
**Phy102 Physics II**  
**Midterm Examination**  
**November 06, 2018 16:30 – 18:30**  
**Good Luck!**

**NAME-SURNAME:**

**SIGNATURE:**

**ID:**

**DEPARTMENT:**


**DURATION:** 120 minutes

- ◇ Answer all the questions.
- ◇ Write the solutions explicitly and clearly.  
Use the physical terminology.
- ◇ You are allowed to use Formulae Sheet.
- ◇ Calculator is allowed.
- ◇ You are not allowed to use any other electronic equipment in the exam.

Question	Grade	Out of
1A		15
1B		15
2		20
3		20
4		20
5		20
<b>TOTAL</b>		110

This page is intentionally left blank. Use the space if needed.

1. A) In Figure, four particles form a square.



The particles have charges  $q_1 = -q_2 = 100 \text{ nC}$  and  $q_3 = -q_4 = 200 \text{ nC}$ , and distance  $a = 5.0 \text{ cm}$ . What are the  $x$  and  $y$  components of the net electrostatic force on particle 3?

$q_1 = 100 \times 10^{-9} \text{ C}$   
 $q_2 = -q_1$   
 $q_3 = 200 \times 10^{-9} \text{ C}$   
 $q_4 = -q_3$   
 $a = 5 \times 10^{-2} \text{ m}$

i)  $F_{3,net,x}$  &  $F_{3,net,y}$ ?  $F_{3,net} = \sum_{i=1}^3 \vec{F}_{2i} = \vec{F}_{31} + \vec{F}_{32} + \vec{F}_{34}$  (2)

$\vec{F}_{3,net,x} = |\vec{F}_{34}| + |\vec{F}_{32}| \cos 45^\circ$  (1)  
 $\vec{F}_{3,net,y} = |\vec{F}_{32}| \sin 45^\circ - |\vec{F}_{31}|$  (1)

$F_{3,net,x} = \frac{k|q_3||q_4|}{a^2} + \frac{k|q_3||q_2|}{(a\sqrt{2})^2} \frac{\sqrt{2}}{2} = \frac{k|q_3|}{a^2} \left( |q_4| + \frac{|q_2|\sqrt{2}}{2} \right) = \frac{8.99 \times 10^9 \text{ Nm}^2/\text{C}^2 (200 \times 10^{-9} \text{ C})}{(5 \times 10^{-2} \text{ m})^2} \left( | -200 \times 10^{-9} \text{ C} | + \frac{| -100 \times 10^{-9} \text{ C} | \sqrt{2}}{2} \right)$   
 $= 0.169 \text{ N}$  (1) (1)

$F_{3,net,y} = \frac{k|q_3|}{a^2} \left( \frac{|q_2|\sqrt{2}}{2} - |q_1| \right) = \frac{8.99 \times 10^9 \text{ Nm}^2/\text{C}^2 (200 \times 10^{-9} \text{ C})}{(5 \times 10^{-2} \text{ m})^2} \left( \frac{100 \times 10^{-9} \text{ C} \sqrt{2}}{2} - 100 \times 10^{-9} \text{ C} \right)$   
 $= -0.046 \text{ N}$  (1) (1)

ii)  $q_1 = q_4 = q$   
 $q_2 = q_3 = -q$   
 $q/q = ?$

$|\vec{F}_{net}| = 0 \rightarrow F_{net,x} = 0$  &  $F_{net,y} = 0$  (1)  
 $|\vec{F}_{net}| = 0 \rightarrow (|\vec{F}_{14}| \cos 45^\circ + |\vec{F}_{12}|) (-\hat{i}) \rightarrow (|\vec{F}_{13}| + |\vec{F}_{14}| \sin 45^\circ) (\hat{j})$

$0 = \frac{k|q_1|}{a^2} \left( \frac{|q_4|\sqrt{2}}{2} + |q_2| \right) = \frac{kq}{a^2} \left( q \frac{\sqrt{2}}{2} + q \right)$  (1)  
 $\Rightarrow \frac{q}{q} = -\frac{4}{\sqrt{2}} = -2\sqrt{2} = -2.83$  (1)

B) In Figure (a), particle 1 (of charge  $q_1$ ) and particle 2 (of charge  $q_2$ ) are fixed in place on an  $x$ -axis,  $8.00 \text{ cm}$  apart. Particle 3 (of charge  $q_3 = +8.00 \times 10^{-19} \text{ C}$ ) is to be placed on the line between particles 1 and 2 so that they produce a net electrostatic force  $F_{3,net}$  on it.






Figure (b) gives the  $x$  component of that force versus the coordinate  $x$  at which particle 3 is placed. The scale of the  $x$  axis is set by  $x_s = 8.0 \text{ cm}$ .

- i What is the sign of charge  $q_1$ ?
- ii What is the ratio  $q_2/q_1$ ?

i)  $\leftarrow \ominus \oplus \ominus \rightarrow$   $\leftarrow F_{31} \quad F_{32} \rightarrow$  ✓ but Figure (b)  
 $\leftarrow \oplus \oplus \oplus \rightarrow$   $\leftarrow F_{32} \quad F_{31} \rightarrow$  ✓ when  $x > 2$  repulsive force (positive value)  
 $\rightarrow q_1$  should be (+)

ii)  $F_{3,net}(x=2) = 0 \rightarrow |F_{32}(x=2)| = |F_{31}(x=2)|$   
 $k \frac{|q_3||q_2|}{(8-x)^2} = k \frac{|q_3||q_1|}{x^2}$  when  $x = 2 \times 10^{-2} \text{ m}$   
 $\frac{q_2}{(6 \times 10^{-2} \text{ m})^2} = \frac{q_1}{(2 \times 10^{-2} \text{ m})^2} \rightarrow \boxed{\frac{q_2}{q_1} = 9}$

2. In the figure below, a nonconducting rod of length  $L = 8.15 \text{ cm}$  has a charge  $q = -4.23 \text{ fC}$  uniformly distributed along its length.



- i) What is the linear charge density of the rod?
- ii) What are the magnitude and direction (relative to the  $+x$ -axis) of the electric field produced at point  $P$ , at distance  $a = 12.0 \text{ cm}$  from the rod?
- iii) What is the electric field magnitude produced at distance  $a = 50.0 \text{ cm}$  by the rod?
- iv) What is the electric field magnitude produced at distance  $a = 50.0 \text{ cm}$  by a particle of charge  $q = -4.23 \text{ fC}$  that replaces the rod?

i)  $\lambda = \frac{q}{L} = \frac{-4.23 \times 10^{-15} \text{ C}}{8.15 \times 10^{-2} \text{ m}} = -5.19 \times 10^{-14} \text{ C/m}$

ii)  $dE = k \frac{dq}{r^2} = k \frac{\lambda dx}{(L+a-x)^2}$   $E = \int_0^L dE$   
 $E_P = k\lambda \int_0^L \frac{dx}{(L+a-x)^2} = k\lambda \left[ \frac{1}{L+a-x} \right]_0^L = k\lambda \left( \frac{1}{a} - \frac{1}{L+a} \right)$   
 $\Rightarrow L = 8.15 \times 10^{-2} \text{ m}$   
 $a = 12 \times 10^{-2} \text{ m}$   
 $E_P = 4.67 \times 10^{-4} \text{ N/C} \left( \frac{8.15 \times 10^{-2} \text{ m}}{(12 \times 10^{-2} \text{ m})(8.15 \times 10^{-2} \text{ m})} \right) = 1.57 \times 10^{-3} \frac{\text{N}}{\text{C}}$

iii)  $L = 8.15 \times 10^{-2} \text{ m}$   
 $a = 50 \times 10^{-2} \text{ m}$   
 $E_P = 4.67 \times 10^{-4} \text{ N/C} \left( \frac{8.15 \times 10^{-2} \text{ m}}{(50 \times 10^{-2} \text{ m})(8.15 \times 10^{-2} \text{ m})} \right) = 1.31 \times 10^{-4} \text{ N/C}$

iv) Point charge:  $E_P = k \frac{|q|}{r^2} = 8.99 \times 10^9 \frac{\text{N}\cdot\text{m}^2}{\text{C}^2} \frac{4.23 \times 10^{-15} \text{ C}}{(50 \times 10^{-2} \text{ m})^2} = 1.54 \times 10^{-4} \text{ N/C}$

3. An infinitely long cylindrical insulating shell of inner radius  $a$  and outer radius  $b$  has a uniform volume charge density  $\rho$ . A line of uniform linear charge density  $\lambda$ , is placed along the axis of the shell. Determine the electric field in the following regions:

i)  $r < a$

ii)  $a < r < b$

iii)  $r > b$

i)  $r < a$

$\oint \vec{E} \cdot d\vec{A} = \frac{Q_{enc}}{\epsilon_0}$   
 $E(2\pi r l) = \frac{Q_{enc}}{\epsilon_0}$   
 $E = \frac{\lambda}{2\pi \epsilon_0 r}$

$Q_{line} = \lambda l$   
 $Q_{cylinder} = \phi$  (shell theorem)  
 $\rightarrow Q_{enc} = \lambda l + \phi$

ii)  $a < r < b$

$\oint \vec{E} \cdot d\vec{A} = \frac{Q_{enc}}{\epsilon_0}$   
 $E(2\pi r l) = \frac{Q_{enc}}{\epsilon_0}$   
 $E = \frac{\lambda + \rho \pi (r^2 - a^2)}{2\pi r}$


$Q_{line} = \lambda l$   
 $Q_{cylinder} = \rho \times \text{Volume}$   
 $= \rho * (\pi r^2 l - \pi a^2 l)$   
 $= \pi l \rho (r^2 - a^2)$   
 $\rightarrow Q_{enc} = \lambda l + \pi l \rho (r^2 - a^2)$

iii)  $r > b$

$\oint \vec{E} \cdot d\vec{A} = \frac{Q_{enc}}{\epsilon_0}$   
 $E(2\pi r l) = \frac{Q_{enc}}{\epsilon_0}$   
 $E = \frac{\lambda + \rho \pi (b^2 - a^2)}{2\pi r}$

$Q_{line} = \lambda l$   
 $Q_{cylinder} = \rho (\pi b^2 l - \pi a^2 l)$   
 $\rightarrow Q_{enc} = \lambda l + \rho l \pi (b^2 - a^2)$

4. Two very thin non-conducting rods are placed together as shown. Both rods have lengths of  $L$  and they carry uniform charges of  $+q$  and  $-q$  over their lengths. Find the potential at point  $P$  at a distance  $a$  and  $d$  from the positively and negatively charged rods as shown. Don't perform integration.




$V_{1 \text{ at } P} = \int dV$   
 $= \int \frac{1}{4\pi\epsilon_0} \frac{dq_1}{r_1}$   
 $dq_1 = \lambda dx$  (3)  
 $r_1 = L + a - x$

$V_{2 \text{ at } P} = \int dV$   
 $= \int \frac{1}{4\pi\epsilon_0} \frac{dq_2}{r_2}$   
 $dq_2 = -\lambda dx$  (3)  
 $r_2 = \sqrt{x^2 + d^2}$

$V = \frac{1}{4\pi\epsilon_0} \frac{q}{r}$   
 $dV = \frac{1}{4\pi\epsilon_0} \frac{dq}{r}$  (2)  
 $dq = ?$   
 $dq = \lambda dx$  (2)

$V_{\text{tot}} = V_{1 \text{ at } P} + V_{2 \text{ at } P} = \frac{\lambda}{4\pi\epsilon_0} \left( \int_0^L \frac{dx}{L+a-x} - \int_0^L \frac{dx}{\sqrt{x^2 + d^2}} \right)$  (3)

5. For the system of capacitors shown in Figure,



find

- i the equivalent capacitance of the system,
- ii the potential across each capacitor,
- iii the charge on each capacitor.

i)  $C_{eq} = ?$

$\frac{3}{3} \parallel \frac{6}{6} \Rightarrow \frac{1}{C_{ac}} = \frac{1}{3\mu F} + \frac{1}{6\mu F} \Rightarrow C_{ac} = 2\mu F$   
 $\frac{2}{2} \parallel \frac{4}{4} \Rightarrow \frac{1}{C_{df}} = \frac{1}{2\mu F} + \frac{1}{4\mu F} \Rightarrow C_{df} = 1.33\mu F$

$\Rightarrow C_{eq} = 3.33\mu F$

ii)

$C = \frac{Q}{V} \sim Q = C_{eq} \times V = (3.33 \times 10^{-6} F) 90V = 299.7\mu C$  (total charge)

$Q_{ac} = (2\mu F) 90V = 180\mu C = q_a = q_c$   
 $Q_{df} = (1.33\mu F) 90V = 119.7\mu C = q_d = q_f$

iii)

$V_a = \frac{q_a}{C_a} = \frac{180\mu C}{3\mu F} = 60V$   
 $V_b = \frac{180\mu C}{6\mu F} = 30V$   
 $V_c = \frac{119.7\mu C}{2\mu F} \approx 60V$   
 $V_d = \frac{119.7\mu C}{4\mu F} \approx 30V$